

Ana Rita de Castro Vieira

Parental Education contributes to a better sleep in the first year of life
A systematic review of the literature

Março, 2018

Ana Rita de Castro Vieira

Parental Education contributes to a better sleep in the first year of life
A systematic review of the literature

Mestrado Integrado em Medicina

Área: Medicina Geral e Familiar

Tipologia: Dissertação

Trabalho efetuado sob a Orientação de:

Prof. Doutor Paulo Alexandre de Azevedo Pereira dos Santos

Trabalho organizado de acordo com as normas da revista:

Prevention Science

Março, 2018

FMUP

Eu, Ana Rita de Castro Vieira, abaixo assinado, nº mecanográfico 199802306, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 13/03/2018

Assinatura conforme cartão de identificação:

Ana Rita de Castro Vieira

NOME

Ana Rita de Castro Vieira

NÚMERO DE ESTUDANTE

199802306

E-MAIL

anacastrovieira@gmail.com

DESIGNAÇÃO DA ÁREA DO PROJECTO

Medicina Geral e Familiar

TÍTULO DISSERTAÇÃO

Parental Education contributes to a better sleep in the first year of life: a systematic review of the literature

ORIENTADOR

Prof. Doutor Paulo Alexandre de Azevedo Pereira dos Santos

COORDINADOR (se aplicável)

ASSINALE APENAS UMA DAS OPÇÕES:

É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTES TRABALHOS APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
É AUTORIZADA A REPRODUÇÃO PARCIAL DESTES TRABALHOS (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTES TRABALHOS.	<input checked="" type="checkbox"/>

Faculdade de Medicina da Universidade do Porto, 19 / 3 / 18

Assinatura conforme cartão de identificação: Ana Rita de Castro Vieira

Agradecimentos

Ao Professor Doutor Paulo Santos por ter aceitado orientar esta dissertação, pela objectividade, rigor científico e inteira disponibilidade.

Aos meus pais, António e Aurora e irmã, Susana, os meus pilares. Obrigada por me incentivarem e apoiarem sempre, por acreditarem mesmo quando eu não acreditei. Foram 6 anos de lutas!

Ao meu cunhado Jonathan, pelo entusiasmo, pelas gargalhadas e amizade.

Aos colegas da turma que me acompanham desde o 1ºano. Não posso deixar de destacar a Ana Sofia, pela paciência e genuína bondade.

Aos meus amigos, Luisa, Emilia, Fiama e Ricardo.

À minha filha Maria Clara, que do alto dos seus 7 anos, responde, muitas vezes, às minhas distrações com um "estudasses!"

Muito obrigada! ...

Parental Education contributes to a better sleep in the first year of life: a systematic review of the literature

Ana Rita Vieira¹, Paulo Santos^{1,2}

¹ Department of Medicine of Community, Information and Health Decision Sciences,
Faculty of Medicine, University of Porto, Portugal

² CINTESIS - Center for Health Technology and Services Research, Faculty of
Medicine, University of Porto, Portugal

Correspondence concerning this article should be addressed to Ana Rita Vieira:
anacastrovieira@gmail.com

Abstract

The first year of life can be overwhelming. One of the major problems parents face is infant's crying and sleeping habits, leading many times to exhaustion, with implications in their sleep, mood, marital satisfaction and mental health. Information is crucial.

The aim was to review the relevance and to assess the impact of parental education in sleep problems in the first year of life. We conducted a systematic review, using keywords reference to "parental education", "parenting", "sleep hygiene", "infant" and "newborn", in Medline via PubMed and Scopus, including Embase, looking for original articles on parental education and infants' sleeping in the first year of life, in the last 10 years in English and Portuguese.

Nine original articles were included in our review. Results suggested that preventive intervention improves infants' sleeping quality. Parental education programs included infant sleep patterns, parental behaviour and strategies facilitating self-soothing. Most successful strategies were placing down the infant while still awake, and to minimize parental responsiveness, by active extinction and graduated extinction techniques. Cognitive-behavioural interventions proved to improve both parents and infants' symptoms.

It is important to include in medical practice parental education on children sleeping pattern and parents' behaviours in children's health routines. It proved to be a cost-effective strategy of increasing empowerment of families to deal with the problem. Formal education of medical doctors is crucial to attend this goal.

Keywords: Parental Education, infant sleep, depression, fatigue, first year of life

Introduction

Sleep problems such as difficulty in falling asleep, frequent night awakenings, difficulty self-soothing and crying are reported by most parents as challenging in the transition to parenthood (Crichton & Symon, 2016).

Inadequate sleep negatively influences infant behaviour, cognitive, physical and socio-emotional development, leading to a diminished ability in emotional regulation (Hall et al., 2015). Fatigue, poor sleep quality and distress are frequent findings. Parents described extreme stages of tiredness with poor parental sleep adding to exhaustion in the first 6 months of life (Loutzenhiser, McAuslan, & Sharpe, 2015).

Sleep problems are present in 30% of infants, depending on how parents define infants' sleep as a problem. Most commonly, parents identify infant night waking with crying beyond six months old as problematic (Meltzer, Plaufcan, Thomas, & Mindell, 2014). Persistent sleep problems can also contribute to postnatal depression, parental suffering and reduced overall health, even in mothers with no depression antecedents (Martin, Hiscock, Hardy, Davey, & Wake, 2007). Moreover, maternal fatigue at 12 months after birth predicted depression scores at 18 months (Giallo, Gartland, Woolhouse, & Brown, 2016). Mothers with higher levels of depression-related symptoms have reported more infant night waking and more maternal worries about adequately responding to their infants' needs at night (Teti & Crosby, 2012). Worries are related with cognitions about infant sleep, particularly doubts about managing infant sleep, difficulty with settling sleep limits and anger at infants' demands around sleep (E. Loutzenhiser, L. Hoffman, & J. Beatch, 2014).

Studies on children's behavioural sleep problems demonstrated the efficacy of behavioural interventions, with significant results that last three to six months after implementation (Mindell et al., 2006). In USA, 14% of mothers experience postnatal

depression (Dave, Petersen, Sherr, & Nazareth, 2010). Similarly, in Portugal, 12,4% of mothers has post-natal depression one week after birth, and 13,7% three months post-partum (Costa, Pacheco, & Figueiredo, 2007). These high rates of maternal depression interfere with the creation of a secure mother-to-child attachment and may cause harmful consequences on the cognitive and emotional growth of the infant (Beck, 1998).

The first year of life is a difficult period of adaptation. Parents need to learn how to deal with the infant's cry and sleep, the two major problems described in the literature. They feel overwhelmed and frequently exhausted and stressed, with repercussions in their own sleep, mood and marital satisfaction. These are the main reasons for them to seek help with their family doctors, paediatricians and assistant nurses (Eckerberg, 2002). Providers must characterize exactly the infant sleep quality, its physiology and development implications (Mindell, Moline, Zendell, Brown, & Fry, 1994).

Although 96% of American paediatricians consider a crucial duty to provide information about infant sleep to the parents, only 18% have formal education in this topic (Faruqui, Khubchandani, Price, Bolyard, & Reddy, 2011). Programs targeting to avoid infant sleep difficulties focus on parental education about baby self-soothing strategies, healthy sleep/wake patterns, little stimulus at night and rising periods among wakening and night feds (B. G. Symon, Marley, Martin, & Norman, 2005). Implementing programs on healthy sleep and crying patterns, and encouraging infants to self-soothe may decrease concerns about infant sleep and crying troubles (Tse & Hall, 2008). However, parents are unsatisfied with counselling obtained from health providers and demand for more information that is acceptable and realistic for practice purposes (Armstrong, Quinn, & Dadds, 1994). However, the way parents

recognize and accomplish behavioural intervention is not totally comprehended (Mindell, 1999).

The literature presents several behaviour strategies helping parents to promoting better infant sleep: positive routines; extinction; gradual or modified extinction and attachment parenting style. Positive routines relates with placing the infant to sleep at consistent hours as same as to wake in the morning, establishing the infant bed as the appropriate place to fall asleep and avoiding associations with behavioural patterns like being rocked, fed or having parents at their bed until they fall asleep. Extinction is a strategy where parents put their infant in bed to sleep in a defined schedule ignoring them until next morning, without checking for potential harm. The method is based in removing the reinforcement for the sleeping problem behaviour. Gradual extinction refers to ignoring infants in their bed for increased periods of time, according to age, temperament of the child and parents' judgement related to the tolerance of their infant to crying. Short interventions of 15 seconds to one minute are allowed, where parents may calm down their child. The aim is to promote infant's ability to self-soothe and fall asleep alone (Kuhn & Elliott, 2003). Extinction-based strategies have been used for half a century, since studies had demonstrated that parental attention reinforce problematic behaviour, diminishing when parents stop attending the child at first demand (Williams, 1959). However, recent studies don't support this technique, showing that the perception of abandoning of the child leads to a great anxiety in parents, and feelings of guilt. Consequently, gradual or modified extinction techniques were introduced to alleviate this guilty perception. Instead of a full extinction program, parents are invited to listen infants cry after a predetermined period, increasing intervals over time (Crichton & Symon, 2016). Attachment parenting style is another opposing method, settled from the values of attachment

theory. It consists on answering rapidly and caringly to the emotional and physical needs of the infant to ease safe attachment (Bowlby, 1978).

There's a prompt need of sleep education programs both in the community or in the primary care settings, as they offer an unique opportunity to improve the insufficient sleep education, with better health and well-being (Gruber, Cassoff, & Knauper, 2011). Working in this gap of parental information leads to increasing awareness and creates formal and systematic knowledge to improve programme design, delivery and effectiveness. Improving sleeping quality leads to better infant health, with lower costs for parents and for health system.

The aim of this review was to assess the impact of parental education in sleep problems in the first year of life, establishing the relevance of this intervention.

Methods

The results of this systematic review are reported according to PRISMA statement (Moher D, Liberati A, Tetzlaff J, & G, 2009).

Search procedure

A search strategy was developed to identify studies correlated to parental education and sleep from birth to the age of 12 months. We searched in Medline via PubMed and Scopus, which includes Embase, for relevant papers published in the last 10 years, whose title, abstract or keywords included reference to “parental education”, “parenting”, “sleep hygiene”, “infant” and “newborn”. The search was limited to English and Portuguese language articles. Titles and abstracts were examined to obtain potential relevant articles.

Exclusion criteria

We excluded non-original articles and those involving premature, infants with biological causes or sleep problems, developmental incapacity or special needs.

Study selection and data extraction

After the systematic search we identified possibly eligible studies and imported them into an Endnote® library. Duplicates were removed. Titles and abstracts were analysed for eligibility and explored full-text articles for potential inclusion. Data were extracted to an excel base. Authors of potentially eligible studies were contacted when needed and reference lists examined for additional articles.

We identified 702 papers (Fig.1) in the first round. After analysing titles and abstracts, we selected 40 articles for full-text reading. The final results included six articles, plus three papers found in references' lists. Not appropriate protocol methods; infant age and unmatched aims with our objectives were the main reasons for exclusion of the articles.




Figure 1 - Selection and exclusion procedure adapted from PRISMA (Mother D et al., 2009)

Quality assessment

The quality was assessed with the Cochrane tool for assessing risk of bias (Higgins & Green, 2011). It was used to rate each study as “low”, “medium” and “high” chance of bias related to selection, performance, detection, attrition and reporting bias.

Results

Database results

This review included six articles from search strategy and three from references' lists.

Study characteristics

We summarized the main aspects of the nine studies included in this review as shown in table one. Articles were published between 2006 and 2016. The majority of studies (n=4) were conducted in Canada (Hall, Clauson, Carty, Janssen, & Saunders, 2006; Hall et al., 2015; E. Loutzenhiser et al., 2014; Tse & Hall, 2008), two were conducted in the USA (Eisenberg et al., 2015; Paul et al., 2016), two in Australia (Hiscock et al., 2007; Hiscock et al., 2014) and one in Japan (Adachi et al., 2009). Most of the trials analysed have one intervention group and one control group. Intervention was delivered orally (group presentation), written (booklet), by video, chart completion and telephone. Themes were about normal infant sleep, bedtime routine, sleep location, behaviour techniques, setting limits, night waking, consequences of insufficient sleep for infants and parents, importance of daytime routines and negative sleep associations. Control groups received no information on infant sleep (Adachi et al., 2009), information about safety (Hall et al., 2015; Paul et al., 2016) or common precaution (Hiscock et al., 2007; Hiscock et al., 2014). We found two qualitative studies about parental perceptions of advice received and effectiveness of a

behavioural intervention (Eisenberg et al., 2015; E. Loutzenhiser et al., 2014).

Globally, the nine studies included 3.312 participants.

Table 1 – Summary of included studies in this systematic review

Study	Age	Participants (N)	Objective	Interventions	Outcomes
(Adachi et al., 2009) - Prospective cohort study, cluster sampling controls - Primary care setting - Japan	4 months	<u>Intervention group:</u> 99 50% male infants 39 gestational weeks Maternal age 32 years 99% married <u>Control group:</u> 95 55% male infants 39 gestational weeks Maternal age 32 years 97% married	Reduction of maternal undesirable behaviours; Reduction of night waking and crying at night.	<u>Intervention group:</u> delivery of a booklet after a health check-up and participation in a brief 10- minute group presentation with guidance on infant sleep. <u>Control group:</u> participants receive no information on infant sleep and provided with only the standard parenting educational information supplied to all visitors at the health check-up session.	Reduction of maternal undesirable behaviours (<i>“feed or check diaper promptly”</i> , <i>“hold and soothe immediately after night waking”</i>) Prevention of increase night waking
(Paul et al., 2016) - Randomized trial - Home visit and health centre - USA	3,16,28 and 40 weeks 1 year (infants)	<u>Intervention group:</u> Responsive parenting (RP) - 140 54% male infants 40 gestational weeks Maternal age 29 years 73% married <u>Control group:</u> 139 50% male infants 40 gestational weeks Maternal 29 years 78% married	Improve sleep behaviours and duration.	<u>Intervention group:</u> messages and guidance within bedtime routine, sleep location and behaviour and night waking (written and video) <u>Control group:</u> information about crib safety and prevention of sudden infant death syndrome, development changes and carbon monoxide poisoning.	Compared with controls, intervention groups were less likely to have prolonged bedtime routines (>45min) and more likely to have earlier bedtimes at 16 and 40 weeks. There were less likely to be fed immediately before bed and more likely to self-soothe to sleep. At 8, 16 and 40 weeks, RP group nocturnal sleep duration was longer by 35, 25 and 22 min, respectively. Sleep duration at 1 year was similar between groups.
(Hall et al., 2015) - Randomized control trial - Home visit and health centre - Canada	6-8 months	<u>Intervention group:</u> 117 64% male infants Parents age 35 years 97% married <u>Control group:</u> 118 48% male infants Parents age 35 years 97% married	Determine the efficacy of a cognitive-behavioural group intervention, offered through public health nurses.	<u>Intervention group:</u> oral and written information about healthy infant sleep, consequences of insufficient sleep for infants and parents, setting limits around infant sleep, the importance of day time and sleep habits and negative sleep connotations. <u>Control group:</u> oral and written information about infant safety risks.	Reduced number of nighty wakes or parent report of sleep problem severity; Improvement in parental depression, fatigue, sleep, and parental cognitions about infant sleep
(Tse & Hall, 2007) - Descriptive, exploratory study - Home visit	6-12 months	14 families mean age: 34 years	Describe parents’ perspectives about implementing sleep intervention, its effectiveness and any problems associated with	Intervention in previous study – EIAR BSP- 2h teaching session with information about infant	Parents altered their evaluations about sleep and parenting styles They experienced many challenges and inadvertent

- Canada			participation in the study.	development, normal sleep behaviour, positive daytime and bedtime routines, negative sleep associations, a controlled comforting intervention and ways which parents could care for themselves.	benefits They welcomed a structured framework for assisting with sleep problems The parents fitted intervention strategies into their realities.
(Eisenberg et al., 2015) - Stratified 2-stage clustered design - Hospitals - USA	2-6 months	1031 mothers 52% male infants 53% of mothers with 20-29 years 37% of mothers with college degree 63% of mothers multiparous	Determine whether mothers report receiving advice from doctors, nurses, family and media. Describe the extent to which advice is consistent with current recommendations.	None (Interviews)	Doctors are the most prevalent source of reported advice; 20% of mothers reported no doctor advice for sleep positioning and more than 50% reported no advice regarding sleep location; Advice from nurses were similar to doctors; Advice from family or media was 20 to 56% for all care practices and advice was often inconsistent with recommendations; Black and Hispanic mothers and first-time mothers were more likely to report recommendation consistent advice.
(L. Loutzenhiser, J. Hoffman, & J. Beatch, 2014) - Internet based study - Canada	6-12 months	411 parents 53% male infants Mean age – 32 years 97% married	Determine parents' use of graduated extinction and their perceptions of its effectiveness	None	Almost half of the parents surveyed reported using graduated extinction with their infants and the majority of these parents began using it before their infants were 6 months old; Parents in the community experienced less success with graduated extinction than parents in clinical setting, with almost half reporting no reduction in infant night waking. Significant predictors of success were parental cognitions regarding stress and support.
(Hall et al., 2006) - Quasi-experimental one group pre-test and post-test design study - Canada	6-12 months	39 families Mean age – 35 years 84,8% have only 1 child 48,8% completing 17-20 years of education	Evaluate the consequences of sleep mediation for infants from 6 to 12 months old	Information about infant sleep and tactics for sleep problems on 2h classes of up to 6 couples, feeding-sleeping and controlled comforting charts completion and bi-weekly telephone call for 2 weeks. Teaching session was about: normal infant sleep; negative	Significant improvement in parents' well being, depressed mood, fatigue and sleep quality. Decreases in parental anger about their infants' sleep and increases in their confidence as parents with consequently more positive interactions with their infants.

				<p>sleep associations; bedtime routines; organized daytime schedules and naps; controlled comforting and take care about themselves (parents).</p> <p>Assessment through questionnaires about parental perceptions of depression, parental sleep quality, sleepiness and fatigue; also cognitions about infant sleep problems and marital adjustment.</p>	
<p>(Hiscock et al., 2007)</p> <ul style="list-style-type: none"> - Cluster randomized trial - Maternal and child health centres - Australia 	7 months	<p>328 mothers</p> <p>89% infant male</p> <p>Maternal age 33 years</p> <p>97% married</p>	Determine whether a community-delivered intervention aiming infant sleep problems improves infant sleep and maternal well-being and to report the costs of this approach to the healthcare system.	<p>Behavioural strategies delivered over individual structured maternal-child health nurses consultations versus usual care.</p>	<p>Prevalence of infant sleep problems was lower in the intervention than control group at 10 months and 12 months. Edinburgh Postnatal Depression Scale (EPDS) scores indicated less depression at 10 months and 12 months</p> <p>SF-12 mental health scores indicated better health at 10 months and 12 months.</p>
<p>(Hiscock et al., 2014)</p> <ul style="list-style-type: none"> - Randomized control trial - Child centres - Australia 	4-6 months	<p>781 infants</p> <p>56% male</p> <p>87% 37-41 weeks of gestation</p> <p>Maternal age 33 years</p> <p>96% married</p>	Evaluate a prevention program for infant sleep problems and postnatal depression.	<p><u>Intervention group:</u> information about healthy infant sleep, cry patterns, soothing methods, biological causes of crying and parent learning of care of them selves, via booklet and DVD, telephone consultation and parent group.</p> <p><u>Control group:</u> well-child care</p>	<p>Intervention caregivers reported a slightly greater reduction in depression symptoms between 4 and 6 months and fewer symptoms at 6 months.</p> <p>Intervention caregivers were less likely to attend night waking, change formulas to manage their infants' problems and had less doubt, difficulty setting limits and concerns about safety with respect to infant sleep.</p> <p>Reductions of sleep problems in infants who are frequent feeders.</p>

Outcomes measured

Primary outcomes described concerns associated with fewer prolonged bedtime routines (>45 min.) and earlier bedtimes, ability to self-soothe, longer nocturnal sleep duration and less parental reports of sleep severity problems.

Secondary outcomes were related with prevention of an increase in night waking, enhancement in parental depression, exhaustion, sleep and perceptions about infant sleep. After intervention, parents altered their perceptions concerning sleep and childrearing styles, using a framework for assisting sleep problems and adapt to their reality. Reduction of maternal undesirable behaviours like *“feed or check diaper promptly”* and *“hold and soothe immediately after night waking”* was also described.

Quality assessment of included studies

We organized the quality assessment chart as shown in table two. Globally, the studies included in this review presented good quality evaluation with low to moderate risk of bias.

Table 2 – Quality assessment of the included studies adapted from Cochrane tool for assessing risk of bias (Higgins & Green, 2011)

Study	Selection bias	Performance bias	Detection bias	Attrition bias	Reporting bias
(Adachi et al., 2009) - Prospective cohort study, cluster sampling controls - Primary care setting Japan	A	B	B	A	A
(Paul et al., 2016) - Randomized control trial - Home visit and health centre USA	A	B	B	A	A
(Hall et al., 2015) - Randomized control trial - Home visit and health centre Canada	B	B	B	A	A
(Tse & Hall, 2007) - Descriptive, exploratory study - Home visit Canada	B	A	A	A	A
(Eisenberg et al., 2015) - Stratified 2-stage clustered design - Hospitals USA	B	A	A	A	A
(L. Loutzenhiser et al., 2014) - Internet based study Canada	B	B	A	A	A

(Hall et al., 2006)	B	B	A	A	A
- Quasi-experimental one group pre-test and post-test design study					
Canada					
(Hiscock et al., 2014)	A	A	B	B	A
- Randomized control trial					
- Child centres					
Australia					
(Hiscock et al., 2007)	A	A	B	A	A
- Cluster randomized trial					
- Maternal and child health centres					
Australia					

Parental Education Programs contribute to better sleep and better parenting

Addachi et al. studied 99 mothers of infants who received a booklet with information about positive sleep routines, desirable and undesirable parental behaviours to place the infant back to sleep after night waking in their routine visit to doctor. The aim was to promote extinction of night waking and improve difficulties in settling to sleep. Desirable strategies included waiting for a short time without responding when the infant cries at night and silently checking bed and clothes of the infant. Undesirable behaviours were the prompt feeding or immediate checking of the infant's diaper after crying as well as prompt holding and soothing. To encourage better sleep, authors recommended increasing daytime play activities, involving abundant visual and hearing stimulation, introducing a regular waking time and bedtime, avoidance of stimulating activities later in the evening, establishing a specific place to sleep at a specific time, setting a bedtime routine and encouraging the infant to fall asleep alone. Intervention group presented less undesirable behaviours, with a significant decrease of *"feed or check diaper promptly"* behaviour from 66,7% to 36,4% and *"hold and soothe immediately"* behaviour from 22,7% to 10,6% versus an increase in control group. There was a meaningful rise of desirable parental behaviours, like *"play with infant or stimulate in the daytime"*, *"settle to sleep at the same place"* and *"set regular bedtime and waking time"*. Generally, the proportion of infants who showed *"difficulty settling"* decreased in both groups (Addachi et al., 2009).

Paul et al. conducted another study, with 270 participants recruited after delivery from one maternity ward. Intervention group was contacted at home at three, four, 16, 18 and 40 weeks, looking for information about infant behavioural states: drowsy, sleeping, fussy and alert. Parents received material by mail, including video

describing alternative soothing strategies, while control group received information just about safety. One year after delivery the evaluation assessed bedtime routine, sleep location and behaviour, night waking and personalized sleep profile. After intervention, more infants were given time to self-soothe to sleep (from 35% to 16%) making them more able to fall asleep alone, without parental intervention. The bedtime routines reduced to less than 45 minutes before bed, infants were less likely to be fed as the last activity of their bedtime routine, less infants felt asleep being held, the self-soothing to sleep increased at 16 and 40 weeks. The intervention group showed also an increase of nighttime sleep duration all over the first year, disappearing after one-year evaluation, although they remained more likely to sleep the recommended 12 to 14 hours per day. Intervention also improved parents' behaviour, reducing the feeding immediately before bedtime at 16 and 40 weeks, and using alternative strategies from picking up and holding/rocking their children for soothing to sleep at 40 weeks (Paul et al., 2016).

Another study assessed the effect of a sleep intervention, including group education sessions with information about infants' patterns, negative sleep associations, unrealistic expectations about sleep, effects of sleep loss on infants and parents, and strategies to reduce night waking. Parents were called by phone twice a week for two weeks. The intervention consisted in a strategy of controlled comforting with two to ten minutes taken to console the infant and then leaving the room for incremental intervals of two minutes up to ten minutes, aiming to promote infant self-settling. Control group was taught just about infant safety protection. The intervention group presented fewer infants' night-time awakening and reported less severe sleep problems (at six weeks, four % in the intervention group compared to 14% of controls). Parents showed significant less fatigue, better sleep quality and

improvement in depressed mood. Also, the doubts about managing infant sleep, comfort managing sleep and feeding, and comfort setting limits around infant sleep improved significantly (Hall et al., 2015).

Eisenberg et al. interviewed 1,031 mothers to know how they get information for caring of their babies. Twenty per cent didn't get any advice from doctors about sleep positioning and beyond 50% stated no guidance concerning secure sleeping places. Advices from nurses were similar to doctors and guidance from family or media was pointed by 20 to 56% for all care practices. It was common to find inconstant advices (Eisenberg et al., 2015).

Loutzenhiser et al. inquired 411 parents. Almost half of them used gradual extinction with their infants, the majority of which starting before six months old. However, parents reported little success as less than half of children got any benefit on night-waking, especially in the community based population, comparing with clinical setting population (E. Loutzenhiser et al., 2014). Parental cognitions regarding stress and support were significant predictors of success (Kuhn & Elliott, 2003).

Hall et al. evaluated consequences of sleep mediation for infants from six to 12 months old in a group of 39 families facing infants' sleep problems. Intervention consisted on delivering material concerning infant sleep and techniques for sleep difficulties on 2h classes of up to six couples, feeding-sleeping and controlled comforting charts completion, and bi-weekly telephone call for two weeks. Teaching session covered the topics about normal infant sleep, bad sleep connotations, bedtime habits, planned daytime plans and naps, organized reassuring, and time available to parents. Assessment was carried out through questionnaires about parental perceptions of depression, parental sleep quality, sleepiness and fatigue, perceptions about infant sleep difficulties and matrimonial adjustment. Authors described

meaningful enhancement in parents' wellbeing, depressed mood, fatigue and sleep quality. They also found a reduction in parental anger about their infants' sleep and an increase in their self-confidence as parents, both associated to more positive interactions with infants (Hall et al., 2006). Two years later, Tse and Hall reassessed 14 of the 35 families who attended the previous intervention, looking for parental perceptions of the specific sleep mediation and any problem linked with participation in the study. Parents mentioned that the information helped to understand the necessity to facilitate their babies to self-soothe, learning that good sleep behaviour and routines leads to a healthier and happier child. Most of the parents read extensively about sleep interventions and a lot of theories about it, but their struggles hadn't accomplished any results while they felt overcome by their babies' cries. Using step-by-step guidelines helped parents to *"gain a framework to tackle sleep problems"*, such as controlled comforting, avoiding feeding to sleep, short naps and co-sleeping, delaying their reply to a partially awake baby or using progress notes to track a baby's pattern of activities. Parents also pointed the *"unanticipated changes resulting from using the strategies in the study"*, related to the notion that they were "normal" parents and their infants' problems were common and modifiable. Thus, they felt empowered for changing their parenting styles, using more routine-oriented schedules rather than totally child-directed approach. Perseverance was the biggest challenge, because parents were asked to refrain from using what they considered easier methods in dealing with crying babies, like cuddling, rocking or breastfeeding to sleep. They identified difficulties attaching to routines, dealing with the effects on others, adapting the intervention to the baby's development, facing with family circumstances that interfered with routines, and dealing with childcare providers who undermine the intervention. Sleep-problems relapses happened with disruptions of

routines, as illnesses, teething or vacations, and they could be very discouraging, bringing the fear of failing in the intervention or being judged. Finally, they mentioned “*parents’ support systems, expectations and inadvertent benefits of the study*” describing varying needs for support. Some of them received help from family members, such as cooking or babysitting, while others received psychological support through phone call or meetings with friends and family, helping to deal with the feeling of being emotionally isolated (Tse & Hall, 2008).

Hiscock et al. conducted a study with 781 babies born at 32 weeks or latter in 42 health-child centres. Intervention families were mailed with a booklet and a DVD containing material concerning healthy infant sleep, crying patterns, soothing methods, crying biological reasons and parents learning to care of them selves. They also received a phone contact at six-eight weeks and participated in parent groups at 12 weeks. Both encouraged parents to discuss crying or sleeping problems and develop problem-solving strategies. The control group received usual health-child care at health services. Intervention caregivers reported a slightly reduction in depression symptoms between four and six months and fewer symptoms at six months. They were less likely to attend night waking, more likely to change formulas to manage their infants’ problems, and they had less doubts and difficulty setting limits and less concerns about safety sleep. The authors reported reduction of sleep problems in infants who wanted to eat frequently at night (Hiscock et al., 2014).

Hiscock et al. studied the impact of a community-delivered intervention aiming baby sleep difficulties, increasing infant sleep pattern, maternal well-being and on the costs to the healthcare system. They followed 328 mothers, between four and 12 months years old, implementing behavioural strategies delivered by maternal-child nurses versus usual care. Mothers could choose between two behavioural methods:

controlled crying, where parents gradually increased the response time to their infant's cry, or camping out, consisting of being seated with the baby till they felt sleeping and progressively eliminating parental presence. Mothers were invited to write sleep diaries to simplify identification of sleep problems and enhancements and to support settlement of further goals. Prevalence of infant sleep difficulties and mother's depression indicators were inferior in the intervention group, at 10 months and 12 months, with better quality of life and acceptable costs, slightly lower in intervention group, although not significant (Hiscock et al., 2007).

Discussion

We found few articles studying the relation between parental education and sleep problems in the first year of life in healthy children.

Our results suggest that active preventive intervention improves sleeping quality in infants, providing education about infant sleep patterns, parental behaviour and strategies to facilitate self-soothing. The key of success relies in direct advice to place the infant in bed while still awake, and to minimize parental responsiveness, based on the efficacy of active extinction and graduated extinction techniques. Moreover, interventions lead to fewer doubts of parents about managing their infants' sleep problems. These findings support the relevance of introducing educational interventions in our preventive approach for infants' health follow-up, although some suggestions that this is a transitory problem, self-extinguishing over time, the neuronal and physiology maturation evolves (Crichton & Symon, 2016; Kuhn & Elliott, 2003; Mindell, 1999; Mindell et al., 2006; B. G. Symon et al., 2005; Williams, 1959). Nevertheless, infants who received behavioural interventions presented more

consistency in improvement of sleeping quality (Gradisar et al., 2016; Mindell et al., 2006).

Our review shows that cognitive-behavioural interventions reduce the number of night awakening and sleeping problems' severity as reported by parents, with less parents' symptoms of depression, fatigue, or sleeping pattern, (Hall, Moynihan, Bhagat, & Wooldridge, 2017; B. Symon, Bammann, Crichton, Lowings, & Tucsok, 2012) and better knowledge about infant sleep. The information can be delivered orally or in written forms, individually or in-group and should be adapted to each case accordingly the providers experience and the population profile (Santos et al., 2014). The distribution of written material for parents has a positive relation with sleep improvement (Gruber et al., 2011) with the advantage of being available in the future, allowing further reviews when effectively needed.

The main strategies for advising are to leave the infant in bed while still awake allowing the self-settlement, to reduce the parental interaction to the minimum, to strengthen the difference between day-time and night-time, and to anticipate the feeding schedule for the end of the afternoon to increase feeding intervals at night.

Although the implementation of educational interventions, both in office visits as in structured group or community actions, is time consuming and consequently costs money, this investment may create value to parents and to health system, being cost-effective. Parents benefit from better understanding of sleeping patterns of their children, promoting the establishment of their own night-time autonomy, with less worries and illness. Health services and providers benefit from that better perception of health status, reducing the number of unnecessary visits, and promoting better satisfaction indexes. However, to be successful in teaching patients, providers must train how to do it, making necessary the introduction of educational programmes for

their own (Santos, Alves, & Simoes, 2017). The need of formal education about sleep programs is crucial to empower health providers for acknowledgment and assessment of sleep troubles, for choosing proper mediations, and for providing an effectively support for families (Gruber et al., 2011).

Conclusion

This systematic review collected the evidence of the relevance of a medical practice towards the improvement of parental education about sleep problems. It reduces undesirable maternal behaviours, making the infant more likely to self-soothe, reducing the number of night awakenings, reducing parental symptoms of depression, fatigue and sleeping disturbances and improving parental cognitions about infant sleeping, within acceptable costs.

Infants' health visits must include questions about sleeping pattern of children and their parents' behaviours, and counselling for the insertion of strategies for improving the perceived problems. Having pre-existing written forms, both in conventional paper and in electronic web-based forms, is a facilitator. Community or group interventions may be useful as they facilitate parenting process and give important peer support.

Compliance with Ethical Standards

Funding

None.

Disclosure of potential conflicts of interest

None.

Ethical approval

None (Not applicable)

Informed Consent

None (Not applicable).

References

- Adachi, Y., Sato, C., Nishino, N., Ohryoji, F., Hayama, J., & Yamagami, T. (2009). A brief parental education for shaping sleep habits in 4-month-old infants. *Clin Med Res*, 7(3), 85-92. doi:10.3121/cmr.2009.814
- Armstrong, K. L., Quinn, R. A., & Dadds, M. R. (1994). The sleep patterns of normal children. *Med J Aust*, 161(3), 202-206.
- Beck, C. T. (1998). The effects of postpartum depression on child development: a meta-analysis. *Arch Psychiatr Nurs*, 12(1), 12-20.
- Bowlby, J. (1978). Attachment theory and its therapeutic implications. *Adolesc Psychiatry*, 6, 5-33.
- Costa, R., Pacheco, A., & Figueiredo, B. (2007). Prevalência e preditores de sintomatologia depressiva após o parto. *Rev. Psiq. Clín*, 34, 157-165.
- Crichton, G. E., & Symon, B. (2016). Behavioral Management of Sleep Problems in Infants Under 6 Months--What Works? *J Dev Behav Pediatr*, 37(2), 164-171. doi:10.1097/DBP.0000000000000257
- Dave, S., Petersen, I., Sherr, L., & Nazareth, I. (2010). Incidence of maternal and paternal depression in primary care: a cohort study using a primary care database. *Arch Pediatr Adolesc Med*, 164(11), 1038-1044. doi:10.1001/archpediatrics.2010.184
- Eckerberg, B. (2002). Treatment of sleep problems in families with small children: is written information enough? *Acta Paediatr*, 91(8), 952-959.
- Eisenberg, S. R., Bair-Merritt, M. H., Colson, E. R., Heeren, T. C., Geller, N. L., & Corwin, M. J. (2015). Maternal Report of Advice Received for Infant Care. *Pediatrics*, 136(2), e315-322. doi:10.1542/peds.2015-0551

Faruqui, F., Khubchandani, J., Price, J. H., Bolyard, D., & Reddy, R. (2011). Sleep disorders in children: a national assessment of primary care pediatrician practices and perceptions. *Pediatrics*, *128*(3), 539-546. doi:10.1542/peds.2011-0344

Giallo, R., Gartland, D., Woolhouse, H., & Brown, S. (2016). "I didn't know it was possible to feel that tired": exploring the bidirectional associations between maternal depressive symptoms and fatigue in a prospective pregnancy cohort study. *Arch Women Mental health*.

Gradisar, M., Jackson, K., Spurrer, N. J., Gibson, J., Whitham, J., Williams, A. S., . . . Kennaway, D. J. (2016). Behavioral Interventions for Infant Sleep Problems: A Randomized Controlled Trial. *Pediatrics*, *137*(6). doi:10.1542/peds.2015-1486

Gruber, R., Cassoff, J., & Knauper, B. (2011). Sleep health education in pediatric community settings: rationale and practical suggestions for incorporating healthy sleep education into pediatric practice. *Pediatr Clin North Am*, *58*(3), 735-754. doi:10.1016/j.pcl.2011.03.006

Hall, W. A., Clauson, M., Carty, E. M., Janssen, P. A., & Saunders, R. A. (2006). Effects on parents of an intervention to resolve infant behavioral sleep problems. *Pediatr Nurs*, *32*(3), 243-250.

Hall, W. A., Hutton, E., Brant, R. F., Collet, J. P., Gregg, K., Saunders, R., . . . Wooldridge, J. (2015). A randomized controlled trial of an intervention for infants' behavioral sleep problems. *BMC Pediatr*, *15*, 181. doi:10.1186/s12887-015-0492-7

Hall, W. A., Moynihan, M., Bhagat, R., & Wooldridge, J. (2017). Relationships between parental sleep quality, fatigue, cognitions about infant sleep, and parental depression pre and post-intervention for infant behavioral sleep problems. *BMC Pregnancy Childbirth*, *17*(1), 104. doi:10.1186/s12884-017-1284-x

Higgins, J., & Green, S. (2011). Cochrane Handbook for Systematic Reviews of Interventions, Version 5.1.0. Retrieved from <http://www.handbook.cochrane.org>

Hiscock, H., Bayer, J., Gold, L., Hampton, A., Ukoumunne, O. C., & Wake, M. (2007). Improving infant sleep and maternal mental health: a cluster randomised trial. *Arch Dis Child*, 92(11), 952-958. doi:10.1136/adc.2006.099812

Hiscock, H., Cook, F., Bayer, J., Le, H. N., Mensah, F., Cann, W., . . . St James-Roberts, I. (2014). Preventing early infant sleep and crying problems and postnatal depression: a randomized trial. *Pediatrics*, 133(2), e346-354. doi:10.1542/peds.2013-1886

Kuhn, B. R., & Elliott, A. J. (2003). Treatment efficacy in behavioral pediatric sleep medicine. *J Psychosom Res*, 54(6), 587-597.

Loutzenhiser, E., Hoffman, L., & Beatch, J. (2014). Parental perceptions of the effectiveness of graduated extinction in reducing infant night-wakings. *Journal of Reproductive and Infant Psychology*, 32:3.

Loutzenhiser, L., Hoffman, J., & Beatch, J. (2014). Parental perceptions of the effectiveness of graduated extinction in reducing infant night-wakings. *Journal of Reproductive and Infant Psychology*, 32:3.

Loutzenhiser, L., McAuslan, P., & Sharpe, D. P. (2015). The trajectory of maternal and paternal fatigue and factors associated with fatigue across the transition to parenthood. *Clinical Psychologist*, 19(1), 15-27. doi:10.1111/cp.12048

Martin, J., Hiscock, H., Hardy, P., Davey, B., & Wake, M. (2007). Adverse associations of infant and child sleep problems and parent health: an Australian population study. *Pediatrics*, 119(5), 947-955. doi:10.1542/peds.2006-2569

Meltzer, L. J., Plaufcan, M. R., Thomas, J. H., & Mindell, J. A. (2014). Sleep problems and sleep disorders in pediatric primary care: treatment recommendations,

persistence, and health care utilization. *J Clin Sleep Med*, 10(4), 421-426.
doi:10.5664/jcsm.3620

Mindell, J. A. (1999). Empirically supported treatments in pediatric psychology: bedtime refusal and night wakings in young children. *J Pediatr Psychol*, 24(6), 465-481.

Mindell, J. A., Kuhn, B., Lewin, D. S., Meltzer, L. J., Sadeh, A., & American Academy of Sleep, M. (2006). Behavioral treatment of bedtime problems and night wakings in infants and young children. *Sleep*, 29(10), 1263-1276.

Mindell, J. A., Moline, M. L., Zendell, S. M., Brown, L. W., & Fry, J. M. (1994). Pediatricians and sleep disorders: training and practice. *Pediatrics*, 94(2 Pt 1), 194-200.

Mother D, Liberati A, Tetzlaff J, & G, A. D. (2009). The PRISMA Group. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 6(7).

Paul, I. M., Savage, J. S., Anzman-Frasca, S., Marini, M. E., Mindell, J. A., & Birch, L. L. (2016). INSIGHT Responsive Parenting Intervention and Infant Sleep. *Pediatrics*, 138(1). doi:10.1542/peds.2016-0762

Santos, P., Alves, L., & Simoes, J. A. (2017). What distinguishes a competent doctor in medical education? *Int J Med Educ*, 8, 270-272. doi:10.5116/ijme.595f.b2ad

Santos, P., Martins, C., Sa, L., Portinha, C., Pessanha, P., Ferreira-Silva, A., . . . Couto, L. (2014). Health education: five years' experience of teaching preventive medicine at the Faculty of Medicine of Oporto University. *Educ Prim Care*, 25(2), 103-107.

- Symon, B., Bammann, M., Crichton, G., Lowings, C., & Tucsok, J. (2012). Reducing postnatal depression, anxiety and stress using an infant sleep intervention. *BMJ Open*, 2(5). doi:10.1136/bmjopen-2012-001662
- Symon, B. G., Marley, J. E., Martin, A. J., & Norman, E. R. (2005). Effect of a consultation teaching behaviour modification on sleep performance in infants: a randomised controlled trial. *Med J Aust*, 182(5), 215-218.
- Teti, D. M., & Crosby, B. (2012). Maternal depressive symptoms, dysfunctional cognitions, and infant night waking: the role of maternal nighttime behavior. *Child Dev*, 83(3), 939-953. doi:10.1111/j.1467-8624.2012.01760.x
- Tse, L., & Hall, W. (2007). A qualitative study os parents' perceptions of a behavioural sleep intervention. *Child: care, health and development*, 34:2.
- Tse, L., & Hall, W. (2008). A qualitative study of parents' perceptions of a behavioural sleep intervention. *Child Care Health Dev*, 34(2), 162-172. doi:10.1111/j.1365-2214.2007.00769.x
- Williams, C. D. (1959). The elimination of tantrum behavior by extinction procedures. *J Abnorm Soc Psychol*, 59, 269.

Normas da Revista - Prevention Science

Inquiries

Editor: Catherine P. Bradshaw, Ph.D., University of Virginia

Inquiries regarding journal policy, manuscript preparation and other such general topics should be sent to the **Managing Editor:**

Sandy Lahn

journal@preventionresearch.org

Manuscript Submission

Manuscripts should be in English and submitted to the Editor's Office via the journal's web-based online manuscript submission and peer-review system at <http://prev.edmgr.com>.

- Acceptable formats for manuscripts are: Word, WordPerfect, RTF, TXT, and LaTeX.
- Acceptable formats for figures (artwork) are: TIFF, GIF, JPEG, EPS, PPT, and Postscript.
- Acceptable formats for supplemental materials are:
 - Text: ASCII, Word, PDF, HTML
 - Tables: Excel, Word, HTML, XHTML, XML
 - Audio and Video: AVI, MPG, Quicktime, RM, MP3, WAV; Animation: GIF, JPEG, Flash/Shockwave
 - Images: GIF, JPEG, TIFF

Publication Policies

Submission is a representation that the manuscript has not been published previously and is not currently under consideration for publication elsewhere.

A statement transferring copyright from the authors (or their employers, if they hold the copyright) to the Society for Prevention Research will be required after the manuscript is accepted for publication. The publisher will supply the necessary forms for this transfer during the publishing process. Such a written transfer of copyright, which previously was assumed to be implicit in the act of submitting a manuscript, is necessary under the U.S. Copyright Law in order for the publisher to carry through the dissemination of research results and reviews as widely and effectively as possible.

Manuscript Preparation

Manuscripts that fail to adhere to the page limits and formatting guidelines herein will be returned without review.

Formatting and Paper Length

The manuscript should be prepared in accordance with the [Publication Manual of the American Psychological Association, Sixth Edition](#), including the citing of references, formatting of tables, figures, metrics and the use of abbreviations, numbers and symbols.

The manuscript should be double-spaced throughout, including the reference section. Please note that we follow the APA recommendation of using a 12-Pt. Times New Roman font.

The paper should not exceed 30 double-spaced typewritten pages INCLUSIVE of title page, abstract, references, tables and figures.

Review your submission for these elements

Order of manuscript pages

- 10 ➤ Title Page with all Author Contact Information & the corresponding author e-mail information.
- 10 ➤ Abstract with keywords
- 10 ➤ Manuscript
- 10 ➤ Compliance with Ethical Standards
- 10 ➤ References
- 10 ➤ Tables and figures

Title Page

A title page on its own page listing the title of the paper and all authors with their academic or professional affiliations is required as the first page of the manuscript.

Abstract and Key Words

An abstract is to be provided on its own page and should contain no more than 250 words. A list of 3-5 descriptive key words (or phrases) is to be provided directly below the abstract. Key words should express the precise content of the manuscript, as they are used for indexing purposes.

Compliance with Ethical Standards section

Authors should include the following just before the References section. (See the section below entitled *Compliance with Ethical Standards* for descriptions and requirements of each):

- 10 ➤ Funding (if applicable)
- 10 ➤ Disclosure of potential conflicts of interest
- 10 ➤ Ethical approval
- 10 ➤ Informed consent

Tables and Figures

Tables and figures should be numbered and referred to by number in the text. Each table/figure should be typed on a separate page, included at the end of the document and should have a descriptive title.

Footnotes

Footnotes should be avoided. When their use is absolutely necessary, footnotes should be numbered consecutively using Arabic numerals and should be typed at the bottom of the page to which they refer. Place a line above the footnote, so that it is set off from the text. Use the appropriate superscript numeral for citation in the text.

Supplemental Materials

Appendices, images, audio and video clips, recordings, lengthy tables and figures that cannot

be included in the main article because of page restrictions can be submitted as supplemental materials. Materials submitted in this manner are included for review during the submission process and are made available to the editors and reviewers. They are also made accessible exactly as submitted from the online version of the published article but are not included with the printed version. Supplemental materials should be referenced in the text by the words “(available online)”.

Page Charges

The journal makes no page charges. Reprints are available to authors, and order forms with the current price schedule are sent with proofs.

Ethical Responsibilities of Authors

This journal is committed to upholding the integrity of the scientific record. As a member of the Committee on Publication Ethics (COPE) the journal will follow the COPE guidelines on how to deal with potential acts of misconduct.

Authors should refrain from misrepresenting research results which could damage the trust in the journal, the professionalism of scientific authorship, and ultimately the entire scientific endeavor. Maintaining integrity of the research and its presentation can be achieved by following the rules of good scientific practice, which include:

- ⑩ ➤ The manuscript has not been submitted to more than one journal for simultaneous consideration.
- ⑩ ➤ The manuscript has not been published previously (partly or in full), unless the new work concerns an expansion of previous work (please provide transparency on the re-use of material to avoid the hint of text-recycling (“self-plagiarism”).
- ⑩ ➤ A single study is not split up into several parts to increase the quantity of submissions and submitted to various journals or to one journal over time (e.g. “salami-publishing”).
- ⑩ ➤ No data have been fabricated or manipulated (including images) to support your conclusions
- ⑩ ➤ No data, text, or theories by others are presented as if they were the author’s own (“plagiarism”). Proper acknowledgements to other works must be given (this includes material that is closely copied (near verbatim), summarized and/or paraphrased), quotation marks are used for verbatim copying of material, and permissions are secured for material that is copyrighted.

Important note: the journal may use software to screen for plagiarism.

- ⑩ ➤ Consent to submit has been received explicitly from all co-authors, as well as from the responsible authorities - tacitly or explicitly - at the institute/organization where the work has been carried out, **before** the work is submitted.
- ⑩ ➤ Authors whose names appear on the submission have contributed sufficiently to the scientific work and therefore share collective responsibility and accountability for the results.

In addition:

- ⑩ ➤ Changes of authorship or in the order of authors are not accepted **after** acceptance of a manuscript.

Requesting to add or delete authors at revision stage, proof stage, or after publication is a serious matter and may be considered when justifiably warranted. Justification for changes in authorship must be compelling and may be considered only after receipt of written approval

from all authors and a convincing, detailed explanation about the role/deletion of the new/deleted author. In case of changes at revision stage, a letter must accompany the revised manuscript. In case of changes after acceptance or publication, the request and documentation must be sent via the Publisher to the Editor-in-Chief. In all cases, further documentation may be required to support your request. The decision on accepting the change rests with the Editor-in-Chief of the journal and maybe turned down. Therefore, authors are strongly advised to ensure the correct author group, corresponding author and order of authors at submission.

Upon request authors should be prepared to send relevant documentation or data in order to verify the validity of the results. This could be in the form of raw data, samples, records, etc.

If there is a suspicion of misconduct, the journal will carry out an investigation following the COPE guidelines. If, after investigation, the allegation seems to raise valid concerns, the accused author will be contacted and given an opportunity to address the issue. If misconduct has been established beyond reasonable doubt, this may result in the Editor-in-Chief's implementation of the following measures, including, but not limited to:

- ⑩ ➤ If the article is still under consideration, it may be rejected and returned to the author.
- ⑩ ➤ If the article has already been published online, depending on the nature and severity of the infraction, either an erratum will be placed with the article or in severe cases complete retraction of the article will occur. The reason must be given in the published erratum or retraction note.
- ⑩ ➤ The author's institution may be informed.

Compliance with Ethical Standards

To ensure objectivity and transparency in research and to ensure that accepted principles of ethical and professional conduct have been followed, authors should include information regarding sources of funding, potential conflicts of interest (financial or non-financial), informed consent if the research involved human participants, and a statement on welfare of animals if the research involved animals.

Authors should include the following statements in a separate section entitled "Compliance with Ethical Standards" before the References when submitting a paper:

- ⑩ ➤ Funding (if applicable)
- ⑩ ➤ Disclosure of potential conflicts of interest
- ⑩ ➤ Research involving Human Participants and/or Animals
- ⑩ ➤ Informed consent

The corresponding author should be prepared to collect documentation of compliance with ethical standards and send if requested during peer review or after publication.

The Editors reserve the right to reject manuscripts that do not comply with the above-mentioned guidelines. The author will be held responsible for false statements or failure to fulfill the above-mentioned guidelines.

Research involving human participants and/or animals

1) Statement of human rights

When reporting studies that involve human participants, authors should include a statement that the studies have been approved by the appropriate institutional and/or national research ethics committee and have been performed in accordance with the ethical standards as laid down in the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards.

If doubt exists whether the research was conducted in accordance with the 1964 Helsinki Declaration or comparable standards, the authors must explain the reasons for their approach, and demonstrate that the independent ethics committee or institutional review board explicitly approved the doubtful aspects of the study.

The following statements should be included in the text before the References section:

Ethical approval: “All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.”

For retrospective studies, please add the following sentence: “For this type of study formal consent is not required.”

2. 2) *Statement on the welfare of animals*

The welfare of animals used for research must be respected. When reporting experiments on animals, authors should indicate whether the international, national, and/or institutional guidelines for the care and use of animals have been followed, and that the studies have been approved by a research ethics committee at the institution or practice at which the studies were conducted (where such a committee exists).

For studies with animals, the following statement should be included in the text before the References section:

Ethical approval: “All applicable international, national, and/or institutional guidelines for the care and use of animals were followed.”

If applicable (where such a committee exists): “All procedures performed in studies involving animals were in accordance with the ethical standards of the institution or practice at which the studies were conducted.”

If articles do not contain studies with human participants or animals by any of the authors, please select one of the following statements:

“This article does not contain any studies with human participants performed by any of the authors.”

“This article does not contain any studies with animals performed by any of the authors.”

“This article does not contain any studies with human participants or animals performed by any of the authors.”

Informed consent

All individuals have individual rights that are not to be infringed. Individual participants in studies have, for example, the right to decide what happens to the (identifiable) personal data gathered, to what they have said during a study or an interview, as well as to any photograph

that was taken. Hence it is important that all participants gave their informed consent in writing prior to inclusion in the study.

Identifying details (names, dates of birth, identity numbers, and other information) of the participants that were studied should not be published in written descriptions, photographs, and genetic profiles unless the information is essential for scientific purposes and the participant (or parent or guardian if the participant is incapable) gave written informed consent for publication. Complete anonymity is difficult to achieve in some cases, and informed consent should be obtained if there is any doubt. For example, masking the eye region in photographs of participants is inadequate protection of anonymity. If identifying characteristics are altered to protect anonymity, such as in genetic profiles, authors should provide assurance that alterations do not distort scientific meaning.

The following statement should be included:

Informed consent: “Informed consent was obtained from all individual participants included in the study.”

If identifying information about participants is available in the article, the following statement should be included:

“Additional informed consent was obtained from all individual participants for whom identifying information is included in this article.”

Disclosure of potential conflicts of interest

Authors must disclose all relationships or interests that could have direct or potential influence or impart bias on the work. Although an author may not feel there is any conflict, disclosure of relationships and interests provides a more complete and transparent process, leading to an accurate and objective assessment of the work. Awareness of real or perceived conflicts of interest is a perspective to which the readers are entitled. This is not meant to imply that a financial relationship with an organization that sponsored the research or compensation received for consultancy work is inappropriate. Examples of potential conflicts of interests **that are directly or indirectly related to the research** may include but are not limited to the following:

- ⑩ ➤ Research grants from funding agencies (please provide the research funder and the grant number)
- ⑩ ➤ Honoraria for speaking at symposia
- ⑩ ➤ Financial support for attending symposia
- ⑩ ➤ Financial support for educational programs
- ⑩ ➤ Employment or consultation
- ⑩ ➤ Support from a project sponsor
- ⑩ ➤ Position on advisory board or board of directors or other type of management relationships
- ⑩ ➤ Multiple affiliations
- ⑩ ➤ Financial relationships, for example, equity ownership or investment interest
- ⑩ ➤ Intellectual property rights (e.g., patents, copyrights and royalties from such rights)
- ⑩ ➤ Holdings of spouse and/or children that may have financial interest in the work

In addition, interests that go beyond financial interests and compensation (non-financial interests) that may be important to readers should be disclosed. These may include but are

not limited to personal relationships or competing interests directly or indirectly tied to this research, or professional interests or personal beliefs that may influence your research.

The corresponding author collects the conflict of interest disclosure forms from all authors. In author collaborations where formal agreements for representation allow it, it is sufficient for the corresponding author to sign the disclosure form on behalf of all authors. Examples of forms can be found here: [COI / Downloads](#)

The following statement should be included if no conflicts of interest exist:

Disclosure of potential conflicts of interest: “The authors declare that they have no conflict of interest.”

After acceptance

Upon acceptance of your article you will receive a link to the special Author Query Application at Springer’s web page where you can sign the Copyright Transfer Statement online and indicate whether you wish to order OpenChoice, offprints, or printing of figures in color.

Once the Author Query Application has been completed, your article will be processed and you will receive the proofs.

Open Choice

In addition to the normal publication process (whereby an article is submitted to the journal and access to that article is granted to customers who have purchased a subscription), Springer provides an alternative publishing option: Springer Open Choice. A Springer Open Choice article receives all the benefits of a regular subscription-based article, but in addition is made available publicly through Springer’s online platform SpringerLink.

[Springer Open Choice](#)

Copyright transfer

Authors will be asked to transfer copyright of the article to the Publisher (or grant the Publisher exclusive publication and dissemination rights). This will ensure the widest possible protection and dissemination of information under copyright laws.

Open Choice articles do not require transfer of copyright as the copyright remains with the author. In opting for open access, the author(s) agree to publish the article under the Creative Commons Attribution License.

Offprints

Offprints can be ordered by the corresponding author.

Color illustrations

Online publication of color illustrations is free of charge. For color in the print version, authors will be expected to make a contribution towards the extra costs.

Proof reading

The purpose of the proof is to check for typesetting or conversion errors and the completeness and accuracy of the text, tables and figures. Substantial changes in content, e.g., new results, corrected

values, title and authorship, are not allowed without the approval of the Editor.

After online publication, further changes can only be made in the form of an Erratum, which will be hyperlinked to the article.